

MIDWEST STONE INSTITUTE

Extracorporeal Shockwave Lithotripsy

In review of the Midwest Stone Institute's ASA/NSAID's list MSI has adopted the following policy with regard to those medications that contain ingredients that may interfere with the blood's ability to clot.

The following medications contain nonsteroidal anti-inflammatory agents. These medications need to be stopped for at least a full 5 days prior to having lithotripsy.

Nonsteroidal Anti-inflammatory Medications:

Addaprin
Advil
Aleve
Anaprox
Anaprox DS
Arthrotec
Cambia
Cataflam
Celebrex
Celecoxib
Clinoril
Co-Advil
Daypro
Diclofenac
Duexis
Dristan Sinus Pain Reliever
Dyspel
EC-Naprosyn
Etodalac
Excedrin IB
Excedrin Aspirin Free
Feldene
Fenoprofen
Flanax
Flector Patch
Flurbiprofen
Genpril
IBU
Ibuprofen
Ibutab

Indocin
Indomethacin
I-Prin
Ketorolac
Ketoprofen
Meclofenamate
Mediproxen
Mefenamic Acid
Meloxicam
Midol Extended Relief
Midol Maximum Strength Cramp Formula
Mobic
Motrin Products
Nabumetone
Nalfon
Naprelan
Naprosyn
Naproxen
Oxaprozin
Pediapain Fever
PennSaid Topical Solution
Piroxicam
Ponstel
Prevacid NapraPAC
Provil
Solaraze Topic Gel
Sprin
Sulindac
Tolmetin
Toradol
Treximet
Vimovo
Voltaren
Zipsor
Zorvolex

The following medications contain aspirin or salicylic acid derivatives. These medications need to be stopped for at least a full 5 days prior to receiving lithotripsy therapy.

Aspirin or Salicylic Medications:

Aceta-Gesic
Acuprin
Adprin

Aggrenox
Alka Seltzer
Amigesic
Anacin
Anodynos
A.P.C.
Argesic
Artha-G
Arthritis Pain Formula
Arthritis Strength Bufferin
Arthrotec
ASA
Ascomp with codeine
Ascriptin
Aspirin products
Aspirin
Azdone
Backache Maximum
Bayer Products
BC powder and tablets
Buffasal
Bufferin
Bufferin Arthritis Strength
Buffex
Buffinol
Butalbital Compound
Butazolidin
Cataflam
Choline Magnesium Trisalicylate
Cope
Damason P
DeWitt's Pain Reliever
Diflunisal
Dipyridamole
Disalcid
Doan's
Durabac
Duraxin
Easprin
Ecotrin
Ed-Flex
Empirin
Empirin with codeine
Equagesic
Equazine-M
Exaprin

Excedrin products
 Extra Strength Anacin
 Fiorinal
 Genprin
 Goody's Extra Strength Headache
 Goody's Back and Body Pain
 Halfprin 81
 Heartline
 Levacet
 Magan
 Magnesium Salicylate
 Marthritic
 Medi-First
 Medi-Seltzer
 Meproamate/Aspirin
 Meprogesic Q
 Midol
 Miniprin
 Mobidin
 Momentum
 MST 600
 No Pain
 Norgesic
 Norwich Extra Strength and Regular
 Strength
 Ohmni-Cen
 Orphenadrine Compound
 Oxycodone and aspirin
 Pabalate-SF
 P-A-C
 Painaid
 Pain Relief Effervescent
 Pamprin Maximum Pain Relief
 Pepto-Bismol
 Percodan
 Percogesic
 Persantine
 Phenylbutazone
 Presalin
 Robomol/ASA
 Roxiprin
 S-A-C
 Saletol
 Salflex
 Salocol
 Salsalate
 Salsitab
 Savaysa (edoxaban)
 Sine-aid IB
 Sodium salicylate
 Sodol
 Stanback Headache Powder
 St. Joseph Aspirin
 Summit
 Synalgos-DC capsules
 Trental
 Tricosal
 Vanquish
 Zorprin

The following medications are anticoagulant medications and need to be stopped for the amount of time listed.

Anticoagulant Medications:

(Consult with cardiologist before discontinuing)

COUMADIN (Need PT/PTT) **OFF 5 days**

WARFARIN (Need PT/PTT) **OFF 5 days**

PLAVIX/Clopidogrel (Check **bleeding time**) **OFF 5 days**

EFFIENT (prasugrel)

PLETAL (Need **Bleeding time**) **OFF 5 days**

HEPARIN (need PT/PTT) **OFF 8 Hours**

LOVENOX (enoxaparin) **OFF 12 Hours**

FRAGMIN (dalteparin) **OFF 24 Hours**

INNOHEP (tinzaparin) **OFF 24 Hours**

ARIXTRA (fondaparinux) **OFF 5 days (Need Bleeding Time)**

ARGATROBAN CHECK PT/PTT
ANGIOMAX(bivalirudin) (CHECK PT/PTT)

PRADAXA (dabigatran) **Off 5 days, Check PTT**

Xarelto (rivaroxaban) **Off 24 hours**

Brilinta (Ticagrelor) **Off 5 days**

Savaysa (edoxaban) **Off 24 hours**

Eliquis (Apixiban) **Off 48 hours**

***Arixtra, Pradaxa, Recludin, Argatroban, Angiomax, Eliquis: Anticoagulant effect prolonged with renal impairment.**

This list is the most common medications prescribed. There are hundreds of other aspirin containing products on the market that may contain generic aspirin. As such, any medication containing aspirin or acetylsalicylic acid should be stopped for 5 days prior to lithotripsy. If you have any questions regarding these medications, please feel free to call us directly.

HERBAL PRODUCTS AND VITAMINS USE SHOULD STOP BEFORE SURGERY Certain vitamins, minerals and "Herbal medications" can affect anesthetic

potency, cause hemodynamic instability, potentiate bleeding, and increase cardiac dysrhythmias. These should be stopped 2 to 3 weeks prior to surgery. (Vitamin E, Echinacea, Ephedra, Feverfew, Garlic, Ginger, Ginkgo biloba, Ginseng, Goldenseal, Kava-kava, Licorice, Saw Palmetto, St. John's Wort, Valerian).

DO NOT STOP TAKING ANY PRESCRIBED MEDICATIONS WITHOUT CHECKING WITH YOUR PHYSICIAN